

# On the Møller Energy-Momentum Complex of the Melvin Magnetic Universe

I. Radinschi\* and I-Ching Yang\*\*

Department of Physics

“Gh. Asachi” Technical University,

Iasi, 6600, Romania and

Department of Natural Science Education,

National Taitung Teachers College, Taitung,

Taiwan 950, Republic of China

iradinsc@phys.tuiasi.ro, \*\*icyang@dirac.phys.ncku.edu.tw

## Abstract

We use the Møller energy-momentum complex to calculate the energy of the Melvin magnetic universe. The energy distribution depends on the magnetic field.

Keywords: Møller energy-momentum complex, Melvin magnetic universe.

PACS: 04. 20.-q, 04. 20.

## Introduction

The subject of energy-momentum localization in general relativity continues to be an open one because there is no given yet a generally accepted expression for the energy-momentum density. Even they are coordinate dependent, various energy-momentum complexes give the same energy distribution for a given space-time.

Aguirregabiria, Chamorro and Virbhadra [1] obtained that the energy-momentum complexes of Einstein [2], Landau and Lifshitz [3], Papapetrou [4], and Weinberg [5] give the same result for the energy distribution for any

Kerr–Schild metric. Also, recently, Virbhadra investigated [6] if these definitions lead to the same result for the most general nonstatic spherically symmetric metric and found they disagree. Only the energy-momentum complex of Einstein gives the same expression for the energy when the calculations are performed in the Kerr–Schild Cartesian and Schwarzschild Cartesian coordinates. The Møller energy-momentum complex allows to compute the energy in any coordinate system.

Some results recently obtained [7]–[10] sustain that the Møller energy-momentum complex is a good tool for obtaining the energy distribution in a given space-time. Also, in his recent paper, Lessner [11] gave his opinion that the Møller definition is a powerful concept of energy and momentum in general relativity. Very important is the Cooperstock [12] hypothesis which states that the energy and momentum are confined to the regions of non-vanishing energy-momentum tensor of the matter and all non-gravitational fields. Also, Chang, Nester and Chen [13] showed that the energy-momentum complexes are actually quasilocal and legitimate expression for the energy-momentum.

In this paper we calculate the energy distribution of the Melvin magnetic universe in the Møller prescription. We use geometrized units ( $G = 1, c = 1$ ) and follow the convention that Latin indices run from 0 to 3.

## Energy in the Møller Prescription

The Melvin magnetic universe [14], [15] is described by the electrovac solution to the Einstein–Maxwell equations and consists in a collection of parallel magnetic lines of forces in equilibrium under their mutual gravitational attraction. The Einstein–Maxwell equations are

$$R_i^k - \frac{1}{2} g_i^k R = 8 \pi T_i^k, \quad (1)$$

$$\frac{1}{\sqrt{-g}} (\sqrt{-g} F^{ik})_{,k} = 4 \pi J^i, \quad (2)$$

$$F_{ij,k} + F_{jk,i} + F_{ki,j} = 0. \quad (3)$$

The energy-momentum tensor of the electromagnetic field is given by

$$T_i^k = \frac{1}{4 \pi} \left[ -F_{im} F^{km} + \frac{1}{4} g_i^k F_{mn} F^{mn} \right]. \quad (4)$$

The electrovac solution corresponds to  $J^i = 0$  and is given by the metric

$$ds^2 = L^2 (dt^2 - dr^2 - r^2 d\theta^2) - L^{-2} r^2 \sin^2 \theta d\varphi^2, \quad (5)$$

where

$$L = 1 + \frac{1}{4} B_0^2 r^2 \sin^2 \theta. \quad (6)$$

The Cartan components of the magnetic field are

$$\begin{aligned} H_r &= L^{-2} B_0 \cos \theta, \\ H_\theta &= -L^{-2} B_0 \sin \theta. \end{aligned} \quad (7)$$

$B_0$  is the magnetic field parameter and is a constant in the solution given by (5) and (6).

The energy-momentum tensor has the non-vanishing components

$$\begin{aligned} T_1^1 &= -T_2^2 = \frac{B_0^2 (1 - 2 \sin^2 \theta)}{8 \pi L^4}, \\ T_0^0 &= -T_3^3 = \frac{B_0^2}{8 \pi L^4}, \\ T_2^1 &= -T_1^2 = \frac{2 B_0^2 \sin \theta \cos \theta}{8 \pi L^4}. \end{aligned} \quad (8)$$

The Møller energy-momentum complex  $M_i^k$  [16] is given by

$$M_i^k = \frac{1}{8 \pi} \chi_i^{kl}, \quad (9)$$

where

$$\chi_i^{kl} = -\chi_i^{lk} = \sqrt{-g} \left( \frac{\partial g_{in}}{\partial x^m} - \frac{\partial g_{im}}{\partial x^n} \right) g^{km} g^{nl}. \quad (10)$$

Also,  $M_i^k$  satisfies the local conservations laws

$$\frac{\partial M_i^k}{\partial x^k} = 0. \quad (11)$$

$M_0^0$  is the energy density and  $M_\alpha^0$  are the momentum density components.

The energy and momentum are given by

$$E = \iiint M_0^0 dx^1 dx^2 dx^3 = \frac{1}{8\pi} \iiint \frac{\partial \chi_0^{0l}}{\partial x^l} dx^1 dx^2 dx^3. \quad (12)$$

For the Melvin magnetic universe we obtain

$$\begin{aligned} \chi_0^{01} &= \frac{B_0^2 r^3 \sin^3 \theta}{(1 + 1/4 B_0^2 r^2 \sin^2 \theta)}, \\ \chi_0^{02} &= \frac{B_0^2 r^2 \cos \theta \sin^2 \theta}{(1 + 1/4 B_0^2 r^2 \sin^2 \theta)}. \end{aligned} \quad (13)$$

After some calculations, applying the Gauss theorem and plugging (13) into (12) we obtain the energy distribution

$$E(r) = \frac{1}{3} B_0^2 r^3 - \frac{1}{15} B_0^4 r^5 + \frac{1}{70} B_0^6 r^7. \quad (14)$$

Put the  $G$  and  $c$  at their places we get

$$E(r) = \frac{1}{3} B_0^2 r^3 - \frac{1}{15} \frac{G}{c^4} B_0^4 r^5 + \frac{1}{70} \frac{G^2}{c^8} B_0^6 r^7. \quad (15)$$

The first term represents twice of the classical value of energy [17] obtained in the Landau and Lifshitz and Papapetrou prescriptions. The other terms are due to the relativistic correction.

## Discussion

Many results recently obtained sustain the viewpoint of Bondi [18]. He gave his opinion that a nonlocalizable form of energy is not admissible in relativity.

We obtain the energy distribution of the Melvin magnetic universe using the energy-momentum complex of Møller. The energy depends on the magnetic field. The result is different as that obtained by Xulu [17] using the energy-momentum complexes of Landau and Lifshitz and those of Papapetrou. The first term represents twice of the classical value of energy [17] obtained in the Landau and Lifshitz and Papapetrou prescriptions. The third term, that is due to the relativistic correction, is twice of value of energy obtained in [17]. Also, the Møller energy-momentum complex does not need to carry out calculations in any particular coordinates.

# 1 Acknowledgments

I. -C. Yang thanks the National Science Council of the Republic of China for financial support under the contract number NSC 90-2112-M-143-003.

## References

- [1] J. M. Aguirregabiria, A. Chamorro and K. S. Virbhadra, *Gen. Rel. Grav.* **25**, 1393 (1996).
- [2] C. Møller, *Ann. Phys. (N.Y.)* **4**, 347 (1958); A. Trautman, in *Gravitation: an Introduction to Current Research*, ed. L. Witten (Wiley, New York, 1962, p.169); R.P. Wallner, *Acta Physica Austriaca* **52**, 121 (1980).
- [3] L.D. Landau and E.M. Lifshitz, *The Classical Theory of Fields* (Pergamon Press, 1987, p.280).
- [4] A. Papapetrou, *Proc. R. Irish. Acad.* **A52**, 11 (1948).
- [5] S. Weinberg, *Gravitation and Cosmology: Principles and Applications of General Theory of Relativity* (John Wiley and Sons, New York, 1972, p.165).
- [6] K. S. Virbhadra, *Phys. Rev.* **D60**, 104041 (1999).
- [7] I-Ching Yang, Wei-Fui Lin and Rue-Ron Hsu, *Chin. J. Phys.* **37**, 113 (1999).
- [8] S.S. Xulu, gr-qc/0010062
- [9] S.S. Xulu, gr-qc/0010068
- [10] I. Radinschi, *Mod. Phys. Lett.* **A16(10)**, 67 (2001).
- [11] G. Lessner, *Gen. Relativ. Gravit.* **28**, 527 (1996).
- [12] F.I. Cooperstock, *Mod. Phys. Lett.* **A14**, 1531 (1999).
- [13] Chia-Chen Chang, J. M. Nester and Chiang-Mei Chen, *Phys. Rev. Lett.* **83**, 1897 (1999).

- [14] M.A. Melvin, Phys. Lett. **8**, 65 (1964).
- [15] F.J. Ernst, J. Math. Phys. **15**, 1409 (1974).
- [16] C. Møller, Ann. Phys. (N.Y.) **4**, 347 (1958).
- [17] S. S. Xulu, Int. J. Mod. Phys. **A15**, 4849 (2000).
- [18] H. Bondi, Proc. R. Soc. London **A427**, 249 (1990).